

CORNERSTONE ASTHMA & ALLERGY ASSOCIATES, LLC

Eric W Bantz, MD and John Y Kim, MD

Diplomates, American Board of Allergy and Immunology

103 Old Marlton Pike Suite 211

Medford, New Jersey 08055

TEL (609) 953-7500 // FAX (609) 953-9085

Antihistamines to Avoid Prior to Allergy Testing

- Actifed
- Advil-PM
- Alavert
- Alaway
- Allegra
- Alka-Seltzer-Allergy
- Astelin
- Astepro
- Atarax
- Benadryl
- Bepreve
- Bromfed
- Cetirizine
- Chlorpheniramine
- Chlortrimeton
- Clarinex
- Claritin
- Clemastine
- Cyproheptadine
- Deconamine
- Desloratidine
- Dimetapp
- Diphenhydramine
- Doxepin
- Doxylamine
- Duradryl
- Dymista
- Epinastine
- Extendryl
- Fexofenadine
- Hydroxyzine
- Ketotifen
- Kronofed-A
- Lastacaft
- Levocetirizine
- Loratadine
- Motrin-PM
- Nyquil
- Olopatadine
- Patanase
- PediaCare-Night Time
- Phenergan
- Promethazine
- Rondec
- Pataday
- Patanol
- Pheniramine
- Rynatan
- Semprex-D
- Tavist
- Triaminic
- Tylenol-PM
- Tylenol-Cold
- Unisom
- Vistaril
- Xyzal
- Zyrtec

Please discontinue all antihistamines for 4 days prior to allergy testing.

Please call us if you have any questions or concerns.